

Noxious Times – Fall 2004, Vol.6, No.3 Issue

Extended Reference List for Pondweed Control Article

California Department of Food and Agriculture survey, 2004.

Eicher, G. 1947. Aniline dye in aquatic weed control. J. Wildlife manage. 11:193-197.

Haller, W. T., J. V. Shireman, and D. F. DuRant. 1980. Fish harvest resulting from mechanical control of *Hydrilla*. Trans. Am. Fish. Soc. 109: 517-520.

Haynes, R. R. 1975. A Revision of North American *Potamogeton* subsection *Pusilli* (Potamogetonaceae). Rhodora 76:564-649.*

Hofstra, D.E. and J.S. Clayton, 2001. Evaluation of Selected Herbicides for the Control of Exotic Submerged Weeds in New Zealand: I. The Use of Endothall, Triclopyr and Dichlobenil, National Institute of Water & Atmospheric Research. Journal of Aquatic Plant Management (2001) 39: 20-24

Legner, E. and T. Fisher. 1980. Impact of *Tilapia zillii* (Gervais) on *Potamogeton pectinatus* L., *Myriophyllum spicatum* var. *exalbescens* Jepson, and mosquito reproduction in lower Colorado Desert irrigation canals. Acta Oecol. Oecologica 1: 3-14.

Lembi, C. A. 1996. Assessment of 2,4-D use in aquatic systems in the United States. In: Biologic and economic assessment of benefits from use of phenoxy herbicides in the United States. NAPIAP Report 1-PA-96. USDA. pp. 179-184.

Lorz, Harold W.; Glenn, Susan W.; Williams, Ronald H.; Kunkel, Clair M.; Norris, Logan A.; Loper, Bobby R. 1979. Effects of selected herbicides on smolting of coho salmon. EPA-600/3-79-071. Corvallis, OR: U.S. Environmental Protection Agency. 103 p.

Madsen J.D., K.D. Getsinger, R.M. Stewart, and C.S. Owens, 2002. Whole Lake Fluridone Treatments for Selective Control of Eurasian Watermilfoil: II. Impacts on Submersed Plant Communities, Minnesota State University. Lake and Reservoir Management (2002) 18(3): 191-200

Madsen, J. D. 2000. Advantages and disadvantages of aquatic plant management techniques. ERDC/EL MP-00-1, U.S. Army Engineer Research and Development Center, Vicksburg, MS.

Mayer, R. J. 1978. Aquatic weed management by benthic semi-barriers. J. Aquat. Plant Manage. 16:31-33.

McCowen, M. C., C. L. Young, S. D. West, S. J. Parra, and W. R. Arnold. 1979. Fluridone a new herbicide for aquatic plant management. J. Aquat. Plant Manage. 17: 27-30.

Nelson, L.S., J.F. Shearer, and M.D. Netherland, 1998. Mesocosm Evaluation of Integrated Fluridone Fungal Pathogen Treatment on Four Submersed Plants, US Army Corps of Engineers. Journal of Aquatic Plant Management (1998) 36(1): 73-77

Netherland, M.D.; J.D. Skogerboe; C.S. Owens; and J.D. Madsen 2000. Influence of water temperature on the efficacy of diquat and endothall versus curlyleaf pondweed. Journal of Aquatic Plant Management 38: 25-32.

Parsons, J.K., K.S. Hamel, J.D. Madsen, and K.D. Getsinger, 2001. The Use of 2,4-D for Selective Control of an Early Infestation of Eurasian Watermilfoil in Loon Lake, Washington, Minnesota State University. Journal of Aquatic Plant Management (2001) 39: 117-125

- Reinert, K.H. and J.H. Rodgers. 1987. Fate and persistence of aquatic herbicides. *Rev. Envtl Contamin Toxicol.* 98:61-98.
- Sanders, D. R. 1981. Biological control technology development. An overview in Proceedings 15th Annual Meeting Aquatic plant Control. Res. Program. Misc. Pap. A-81-3. U.S. Army Corps Engineers, Vicksburg, MS. 139-143.
- SePRO Corporation, 11550 North Meridian Street, Suite 600, Carmel, IN 46032 USA.
- Serns, S. L. 1977. Effects of dipotassium endothall on rooted aquatics and adult and first generation bluegills. *Water Res. Bull.* 13(1): 71-80.
- Simsiman, GV., T.C. Daniel and G. Chesters. 1976. Diquat and endothall: Their fates in the environment. *Residue reviews.* 62: 131-174.
- Sisneros, D., M. Lichwardt, and T. Greene. (1998) Low-dose metering of Endothall for aquatic plant control in flowing water., *Journal of Aquatic Plant Management* 36: 69-72
- Spencer, D. F. and Ksander, G. G. 1997. Dilute acetic acid exposure enhances electrolyte leakage by *Hydrilla verticillata* and *Potamogeton pectinatus* tubers. *Journal of Aquatic Plant Management* 35: 25-30.
- Spencer, D. F., C. L. Elmore, G. G. Ksander, and J. A. Roncoroni. Influence of dilute acetic acid treatments on American pondweed winterbuds in the Nevada Irrigation District, California.2003. *Journal of Aquatic Plant Management* 41: 65-68.
- Sprecher, S. L., K. D. Getsinger, A. B. Stewart. 1998. Selective effects of aquatic herbicides on sago pondweed. *J. Aquat. Plant Manage.* 36: 64-68.
- Steward, K. K. 1980. Retardation of hydrilla (*Hydrilla verticillata*) regrowth through chemical control of vegetative propagules. *Weed Sci.* 28 (3):245-251.
- Sutton, D. L., W. T. Haller, K. K. Steward and R. D. Blackburn. 1972. Effect of copper on uptake of diquat 14C by hydrilla. *Weed Sci.* 20(6): 581-583.
- Sutton, D. L., L. W. Weldon and R. D. Blackburn. 1970. Effect of diquat on uptake of copper in aquatic plants. *Weed Sci.* 18(6): 703-707.
- Walker, C. R. 1964. Dichlorobenil as a herbicide in fish habitats. *Weeds.* 12:267-269.
- Wells, R. D. S. and J. S. Clayton. 1993. Evaluation of endothall for aquatic weed control in New Zealand. Proc. 46th NZ Plant Protection Conf. pp. 102-106.
- Westerdahl, H. E., and Getsinger, K. D., eds. (1988). "Aquatic plant identification and herbicide use guide; Volume I: Aquatic herbicides and application equipment," Technical Report A-88-9, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS. NTIS No. AD A195 571.
- Yeo, R. R., and N. Dechoretz. 1977. Acute toxicity of a herbicidal combination of diquat, plus copper ion to eggs, alewife and fry of rainbow trout and to aquatic invertebrates. *J. Aquat. Plant Manage.* 14:57-60.